

Chapter 7

Exercise 7.1

- (a) $H \neg p$
 p : It is warm
- (b) $F p$
 p : There is a sea battle
- (c) $\neg F P p$ or, perhaps, $F \neg P p$
 p : There is a sea battle
- (d) $F(p \vee P q)$
 p : It is warm
- (e) $\neg P p \rightarrow \neg F q$, or $G(\neg P p \rightarrow \neg q)$
 p : You study, q : you pass the exam
- (f) $P(p \wedge q)$
 p : I am having tea, q : the door bell rings

Exercise 7.2

(a), (c), (f), (g), and (h) are true, (b), (d), and (e) are false.

Exercise 7.3

(Con1): Suppose some sentence of the form $A \rightarrow G P A$ is false at some time t in some temporal model. By clause (e) of definition 7.2, this means that A is true at t and $G P A$ is false at t . By clause (h), the latter means that there is a time s with $t < s$ such that $P A$ is not true at s . By clause (i), it follows that A is not true at t . Contradiction.

The argument for (Con2) is analogous.

Exercise 7.4

8 Answers to the Exercises

- (a) 1. $\neg(A \rightarrow GPA)$ (t) (Ass.)
2. A (t) (1)
3. $\neg GPA$ (t) (1)
4. $t < s$ (3)
5. $\neg PA$ (s) (3)
6. $\neg A$ (t) (4,5)
x
- (b) 1. $\neg(A \rightarrow HFA)$ (t) (Ass.)
2. A (t) (1)
3. $\neg HFA$ (t) (1)
4. $s < t$ (3)
5. $\neg FA$ (s) (3)
6. $\neg A$ (t) (4,5)
x
- (c) 1. $\neg(FA \rightarrow HFFA)$ (t) (Ass.)
2. FA (t) (1)
3. $\neg HFFA$ (t) (1)
4. $s < t$ (3)
5. $\neg FFA$ (s) (3)
6. $\neg FA$ (t) (4,5)
x

(d)	1.	$\neg(HA \leftrightarrow HFHA)$	(t)	(Ass.)	
		\swarrow			
	2.	HA	(t)	(1)	
	3.	$\neg HFHA$	(t)	(1)	
	6.	$s < t$		(3)	
	7.	$\neg FHA$	(s)	(3)	
	8.	$\neg HA$	(t)	(6,7)	
		x			
					\searrow
	4.	$\neg HA$	(t)	(1)	
	5.	HFHA	(t)	(1)	
	9.	$s < t$		(4)	
	15.	$\neg A$	(s)	(4)	
	11.	FHA	(s)	(5,9)	
	12.	$s < r$		(11)	
	13.	HA	(r)	(11)	
	14.	A	(s)	(12,13)	
		x			

Exercise 7.5

Suppose $<$ is transitive, and $x > y$ and $y > z$. Equivalently, $y < x$ and $z < y$. By transitivity of $<$, we have $z < x$. So $x > z$.

Exercise 7.6

Suppose R is transitive. If there are points x and y for which xRy and yRx then xRx by transitivity. So if R isn't asymmetric then it isn't irreflexive. If R isn't irreflexive then there is a point x with xRx . This violates asymmetry, because asymmetry demands that if xRx then not xRx .

Exercise 7.7

If time is transitive and circular, then it is neither asymmetric nor irreflexive.

Exercise 7.8

(a), (b), (e), and (f) are invalid. Trees for (c), (d), and (g):

8 Answers to the Exercises

- (c) 1. $\neg(\text{P G } p \rightarrow \text{P F } p)$ (t) (Ass.)
2. $\text{P G } p$ (t) (1)
3. $\neg \text{P F } p$ (t) (1)
4. $s < t$ (2)
5. $\text{G } p$ (s) (2)
6. p (t) (4,5)
7. $\neg \text{F } p$ (s) (3,4)
8. $\neg p$ (t) (4,7)
x
- (d) 1. $\neg(\text{P G G } p \rightarrow \text{G G } p)$ (t) (Ass.)
2. $\text{P G G } p$ (t) (1)
3. $\neg \text{G G } p$ (t) (1)
4. $s < t$ (2)
5. $\text{G G } p$ (s) (2)
6. $t < r$ (3)
7. $\neg \text{G } p$ (r) (3)
8. $s < r$ (3,6,Trans.)
9. $\text{G } p$ (r) (5,8)
x

(g)	1. $\neg(\mathbf{F}(\mathbf{G}q \wedge \neg p) \rightarrow \mathbf{G}(p \rightarrow (\mathbf{G}p \rightarrow q)))$	(t) (Ass.)
	2. $\mathbf{F}(\mathbf{G}q \wedge \neg p)$	(t) (1)
	3. $\neg \mathbf{G}(p \rightarrow (\mathbf{G}p \rightarrow q))$	(t) (1)
	4. $t < s$	(2)
	5. $\mathbf{G}q \wedge \neg p$	(s) (2)
	6. $\mathbf{G}q$	(s) (5)
	7. $\neg p$	(s) (5)
	8. $t < r$	(3)
	9. $\neg(p \rightarrow (\mathbf{G}p \rightarrow q))$	(r) (3)
	10. p	(r) (9)
	11. $\neg(\mathbf{G}p \rightarrow q)$	(r) (9)
	12. $\mathbf{G}p$	(r) (11)
	13. $\neg q$	(r) (11)
	14. $s < r$	
	15. $s = r$	
	16. $r < s$	
	17. q	(r) (6,14)
	18. p	(s) (10,15)
	19. p	(s) (12,16)
	x	x
	x	x

Exercise 7.9

- (a) For example, $\mathbf{G}A \rightarrow \mathbf{F}A$.
- (b) For example, $\mathbf{H}A \rightarrow \mathbf{P}A$.
- (c) No schema corresponds to the class of frames with a last time. If we also assume transitivity and quasi-connected (see page 149), then $\mathbf{G}(A \wedge \neg A) \vee \mathbf{F}\mathbf{G}(A \wedge \neg A)$ works.
- (d) No schema corresponds to the class of frames with a first time. If we also assume transitivity and quasi-connectedness, then $\mathbf{H}(A \wedge \neg A) \vee \mathbf{P}\mathbf{H}(A \wedge \neg A)$ works.

Exercise 7.10

Assume a frame is dense. Suppose for reductio that some instance of $\mathbf{F}A \rightarrow \mathbf{F}\mathbf{F}A$ is false at some point t in some model M based on that frame. Then $\mathbf{F}A$ is true at t and

$\text{FF}A$ is false. Since $\text{F}A$ is true at t , it follows by definition 7.2 that A is true at some point s such that $t < s$. By density, there is a point r such that $t < r < s$. But since A is true at s , $\text{F}A$ is true at r , and so $\text{FF}A$ is true at t ; contradiction.

In the other direction, we have to show that if a frame isn't dense then some instance of $\text{F}A \rightarrow \text{FF}A$ is false at some point t in some model M based on that frame. We take the simplest instance $\text{F}p \rightarrow \text{FF}p$. If a frame isn't dense then there are points t, s such that $t < s$ and no point lies in between t and s . Let V be an interpretation function that makes p true at s and false everywhere else. Then $\text{F}p$ is true at t but $\text{FF}p$ is false. So $\text{F}p \rightarrow \text{FF}p$ is false at t .

Exercise 7.11

Without assumptions about the flow of time there is no way to express in \mathfrak{L}_T that p is true at all times (or at some time). In linear flows, $p \wedge \text{H}p \wedge \text{tG}p$ does the job.

Exercise 7.12

Suppose some instance of $\text{F}A$ is true at the present time t . Then $\text{HFF}A$ is true at t as well. Since $\text{HFF}A$ is equivalent to $\neg \text{P} \neg \text{F}A$, it follows by (S2) that $\Box \text{HFF}A$ is true at t . Since t is not the first point in time, we also have $\text{P}(A \vee \neg A)$ at t . By (S1), we get $\Box \text{P}(A \vee \neg A)$. Now $\text{HFF}A$ and $\text{P}(A \vee \neg A)$ together K_t -entail $\text{F}A$. Assuming that the box is closed under logical consequence, we can infer that $\Box \text{F}A$ is true at t .

Exercise 7.13

Consider a model with three times ordered by $s < t$ and $s < r$. Assume p is true at t and not at r . Then $p \rightarrow \text{HF}p$ is false on the Peircean interpretation.

Exercise 7.14

(a)–(d) are valid, (e) is invalid.

To show that a schema is valid, assume for reductio that there is some time t on some history H in some model M at which the schema is false. Then (repeatedly) use definition 7.3 to derive a contradiction.

For (e), consider a model with three times t, s, r such that $s < t, r < t$, and neither $s < r$ nor $r < s$. Let q be true at s and false at the other two times. $\text{P}q \rightarrow \Box \text{P} \Diamond q$ is

false at t on the history $\langle s, t \rangle$.

Exercise 7.15

A sentence A is super-valid iff $M, t \models A$ for all temporal models M and times t in M . By supervaluationism, this holds iff $M, H, t \models A$ for all M, t , and histories H containing t . That's how Ockhamist validity was originally defined.

Exercise 7.16

$(A \wedge \neg A) \cup A$.

Exercise 7.17

$A p \rightarrow p$.